

COUNTER EXAMPLE TO A STRONG MATROID MINOR CONJECTURE

SHRAWAN KUMAR

Abstract: The main result of this note asserts that a strong form of the Matroid Minor Conjecture due to J. Draisma is not true, i.e., there exist properly ascending chains of S_∞ -stable ideals in the affine coordinate ring of the *affine infinite Grassmannian*, where S_∞ is the infinite symmetric group. In fact, we explicitly construct such an ascending chain. His conjectures on topological noetherian property for the affine infinite Grassmannian remain open though.

1. INTRODUCTION

Let k be any (possibly finite) field. For positive integers n, p , let $V_{n,p}$ be the vector space over k with basis

$$\{x_{-n}, x_{-n+1}, \dots, x_{-1}, x_1, x_2, \dots, x_p\}.$$

Let $\{x_{-n}^*, x_{-n+1}^*, \dots, x_{-1}^*, x_1^*, x_2^*, \dots, x_p^*\}$ be the dual basis of the dual vector space $V_{n,p}^*$, i. e., $x_j^*(x_k) = \delta_{j,k}$.

Let $\text{Gr}(p, V_{n,p}^*)$ be the Grassmannian of p -planes in $V_{n,p}^*$. Then, we have the Plücker embedding:

$$\iota : \text{Gr}(p, V_{n,p}^*) \hookrightarrow \mathbb{P}(\wedge^p(V_{n,p}^*)), \quad A \mapsto \wedge^p(A), \quad \text{for } A \in \text{Gr}(p, V_{n,p}^*).$$

Let $\widetilde{\text{Gr}}(p, V_{n,p}^*)$ be the corresponding affine cone (under the above embedding). Thus, we get an embedding

$$\widetilde{\text{Gr}}(p, V_{n,p}^*) \hookrightarrow \wedge^p(V_{n,p}^*).$$

This makes $\widetilde{\text{Gr}}(p, V_{n,p}^*)$ a closed irreducible (affine) subvariety of $\wedge^p(V_{n,p}^*)$.

Define the surjective maps

$$\xi = \xi^{n+1,p} : \wedge^p(V_{n+1,p}^*) \twoheadrightarrow \wedge^p(V_{n,p}^*)$$

induced from the standard inclusion of the bases and

$$\theta = \theta^{n,p+1} : \wedge^{p+1}(V_{n,p+1}^*) \twoheadrightarrow \wedge^p(V_{n,p}^*), \quad \omega \mapsto i_{x_{p+1}}\omega,$$

where i is the interior multiplication defined by

$$(i_{x_{p+1}}\omega)(v_1, \dots, v_p) = \omega(x_{p+1}, v_1, \dots, v_p), \quad \text{for } v_i \in V_{n,p}.$$

Now, the maps $\theta^{n+1,p+1}$ and $\xi^{n+1,p+1}$ induce the restriction maps which are surjective:

$$\widetilde{\text{Gr}}(p, V_{n+1,p}^*) \xleftarrow{\widetilde{\theta}} \widetilde{\text{Gr}}(p+1, V_{n+1,p+1}^*) \xrightarrow{\widetilde{\xi}} \widetilde{\text{Gr}}(p+1, V_{n,p+1}^*).$$

Define a partial order \leq on $\mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$ by $(n, p) \leq (m, q)$ if $n \leq m$ and $p \leq q$. The above maps $\widetilde{\theta}$ and $\widetilde{\xi}$ give rise to a surjective map between the affine varieties (see §2 for more details):

$$\widetilde{\text{Gr}}(q, V_{m,q}^*) \rightarrow \widetilde{\text{Gr}}(p, V_{n,p}^*) \quad \text{for } (n, p) \leq (m, q).$$

Define the affine schemes:

$$\widetilde{\text{Gr}}(\infty/2, V_\infty^*) := \varprojlim_{(n,p)} \widetilde{\text{Gr}}(p, V_{n,p}^*), \quad \text{and} \quad \wedge^{\infty/2}(V_\infty^*) := \varprojlim_{(n,p)} \wedge^p(V_{n,p}^*).$$

Define the group

$$\mathrm{GL}(\infty) := \varinjlim_{(n,p)} \mathrm{GL}(V_{n,p}), \quad \text{and its subgroup } \mathrm{N}(\infty) := \varinjlim_{(n,p)} \mathrm{N}(V_{n,p}),$$

where $\mathrm{N}(V_{n,p})$ denotes the normalizer of the standard maximal torus in $\mathrm{SL}(V_{n,p})$. The standard action of $\mathrm{GL}(V_{n,p})$ on $\wedge^p(V_{n,p}^*)$ gives rise to an action of $\mathrm{GL}(\infty)$ on $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^*)$ (cf. §2 for more details). Define the infinite symmetric group

$$S_\infty := \varinjlim_n S_n, \quad \text{where } S_n \text{ is the symmetric group on the symbols } \{-n, -(n-1), \dots, -1, 1, 2, \dots, n\}.$$

Then, S_∞ canonically embeds in $\mathrm{GL}(\infty)$ via the permutation matrices.

Jan Draisma kindly told us his following conjecture.

Conjecture 1.1. *Let k be any field. The affine infinite Grassmannian $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^*)$ is topologically S_∞ -noetherian, i.e., every descending chain of S_∞ -stable Zariski-closed subsets of the k -points of $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^*)$ stabilizes.*

A slightly weaker form of the conjecture states that $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^)$ is topologically $\mathrm{N}(\infty)$ -noetherian.*

A stronger form of the conjecture states that any ascending chain of S_∞ -stable ideals in the affine coordinate ring of $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^)$ stabilizes.*

The main result of this note is the following (cf. Theorem 3.4).

Theorem 1.2. *The strong form of the above conjecture is false, i.e., there exists an ascending chain of S_∞ -stable ideals in the affine coordinate ring of $\widetilde{\mathrm{Gr}}(\infty/2, V_\infty^*)$ which does not stabilize.*

In fact, we give such an example explicitly.

Before we can explain the significance of Draisma's Conjecture 2.6 to some important results in Graph Theory, we need to briefly explain some of the very significant results in *Matroid Minors Theory*.

We begin by recalling the following conjecture due to Rota [Ro].

Conjecture 1.3. *For each finite field k , there are, up to isomorphism, only finitely many excluded minors for the class of F -representable matroids.*

Rota's conjecture is reminiscent of the classical Generalized Kuratowski Theorem [Kur]. As part of the *Graph Minors Project*, Neil Robertson and Paul Seymour were able to further generalize the Generalized Kuratowski Theorem to obtain the WQO (*Well Quasi Ordering*) Theorem stated below (cf. [RS]). Their results were published in a series of twenty three journal papers totaling more than 700 pages from 1983 to 2004. Diestel, in his book on graph theory [Di], says that this theorem dwarfs any other result in graph theory and may doubtless be counted among the deepest theorems that mathematics has to offer.

Theorem 1.4. (*WQO*). *Each minor-closed class of graphs has only finitely many excluded minors.*

Equivalently, in any infinite set S of graphs, there must be a pair of graphs one of which is a minor of the other.

Then, Robertson and Seymour proposed ideas for extending their Graph Minors Project to matroids. The challenge was taken up by Jim Geelen, Bert Gerards, and Geoff Whittle. Though it is not true that the WQO Theorem extends to all matroids. However, after extensive work for several years, they (Geelen et al.) announced the following slightly weaker theorem (cf. [GGW, Theorem 6]) significantly extending the WQO theorem to matroids.

Theorem 1.5. (*Matroid WQO Theorem*). *For each finite field k and each minor-closed class of k -representable matroids, there are only finitely many k -representable excluded minors.*

According to their article [GGW], to quote them: ‘We are now immersed in the lengthy task of writing up our results. Since that process will take a few years, we have written this article offering a high-level preview of the proof.’

The following result (communicated to us by J. Draisma), which is fairly easy to prove, provides a direct bridge between Conjecture 2.6 and Theorem 1.5 once we observe that $\theta^{n,p+1} : \wedge^{p+1}(V_{n,p+1}^*) \rightarrow \wedge^p(V_{n,p}^*)$ and $\xi^{n+1,p} : \wedge^p(V_{n+1,p}^*) \rightarrow \wedge^p(V_{n,p}^*)$, at the level of matroids, correspond to contraction and deletion respectively.

Theorem 1.6. *Let k be a finite field and assume that the set of k -points of the affine infinite Grassmannian $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$, equipped with the Zariski topology, is S_∞ -noetherian. Then, matroids representable over k are well quasi ordered by the minor order.*

Acknowledgements: We are indebted to Jan Draisma for explaining to us his Conjecture 2.6, showing its significance via his Theorem 1.6 to the Matroid WQO Theorem 1.5, providing all the references [Di], [GGW], [Kur], [RS], [Ro]. We also thank Andrew Snowden for his comment (see Theorem 3.4). This work was completed while the author was visiting the Institut des Hautes Études Scientifiques (Bures-sur-Yvette, France) during the fall semester of 2023, hospitality of which is gratefully acknowledged. We thank the Referee for some helpful comments.

2. INFINITE GRASSMANNIAN AND THE MATROID MINOR CONJECTURE

The base field in this note is any (possibly finite) field.

Definition 2.1. For positive integers n, p , let $V_{n,p}$ be the vector space over k with basis

$$\{x_{-n}, x_{-n+1}, \dots, x_{-1}, x_1, x_2, \dots, x_p\}.$$

Let $\{x_{-n}^*, x_{-n+1}^*, \dots, x_{-1}^*, x_1^*, x_2^*, \dots, x_p^*\}$ be the dual basis of the dual vector space $V_{n,p}^*$.

Define the linear maps

$$\eta = \eta^{n+1,p} : \wedge^p(V_{n,p}) \hookrightarrow \wedge^p(V_{n+1,p})$$

induced from the standard inclusion of the bases and

$$\beta = \beta^{n,p+1} : \wedge^p(V_{n,p}) \hookrightarrow \wedge^{p+1}(V_{n,p+1}), \quad \omega \mapsto \omega \wedge x_{p+1}.$$

Dually, we get surjective maps

$$\xi = \xi^{n+1,p} : \wedge^p(V_{n+1,p}^*) \twoheadrightarrow \wedge^p(V_{n,p}^*)$$

and

$$\theta = \theta^{n,p+1} : \wedge^{p+1}(V_{n,p+1}^*) \twoheadrightarrow \wedge^p(V_{n,p}^*), \quad \omega \mapsto i_{x_{p+1}} \omega,$$

where i is the interior multiplication. Observe that in Matroid theory, the map ξ corresponds to *deletion* and the map θ corresponds to *contraction*.

Let $\text{Gr}(p, V_{n,p}^*)$ be the Grassmannian of p -planes in $V_{n,p}^*$ (For generalities on Grassmannians, see [EH, §III.2.7].). Then, we have the Plücker embedding:

$$\iota : \text{Gr}(p, V_{n,p}^*) \hookrightarrow \mathbb{P}(\wedge^p(V_{n,p}^*)), \quad A \mapsto \wedge^p(A), \quad \text{for } A \in \text{Gr}(p, V_{n,p}^*).$$

Let $\widetilde{\text{Gr}}(p, V_{n,p}^*)$ be the corresponding affine cone (under the above embedding). Thus, we get an embedding

$$\widetilde{\text{Gr}}(p, V_{n,p}^*) \hookrightarrow \wedge^p(V_{n,p}^*),$$

where the image consists of the decomposable vectors (including the vector 0). This makes $\widetilde{\text{Gr}}(p, V_{n,p}^*)$ a closed irreducible (affine) subvariety of $\wedge^p(V_{n,p}^*)$. Now, the maps

$$(1) \quad \wedge^p(V_{n+1,p}^*) \xleftarrow{\theta^{n+1,p+1}} \wedge^{p+1}(V_{n+1,p+1}^*) \xrightarrow{\xi^{n+1,p+1}} \wedge^{p+1}(V_{n,p+1}^*)$$

induce the restriction maps which are surjective:

$$(2) \quad \widetilde{\text{Gr}}(p, V_{n+1,p}^*) \xleftarrow{\bar{\theta}} \widetilde{\text{Gr}}(p+1, V_{n+1,p+1}^*) \xrightarrow{\bar{\xi}} \widetilde{\text{Gr}}(p+1, V_{n,p+1}^*).$$

To prove the existence of $\bar{\theta}$, we can write

$$v_1 \wedge \cdots \wedge v_{p+1} = v'_1 \wedge \cdots \wedge v'_p \wedge (v'_{p+1} + \alpha x_{p+1}^*), \text{ where } v'_i(x_{p+1}) = 0 \forall 1 \leq i \leq p+1,$$

for some $\alpha \in k$. Thus,

$$i_{x_{p+1}}(v_1 \wedge \cdots \wedge v_{p+1}) = \pm \alpha v'_1 \wedge \cdots \wedge v'_p$$

and hence θ induces the map $\bar{\theta}$ on the corresponding cones of Grassmannians. The existence of $\bar{\xi}$ is trivial to see.

Define a partial order \leq on $\mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$ by

$$(3) \quad (n, p) \leq (m, q) \text{ if } n \leq m \text{ and } p \leq q.$$

We have a surjective map between the affine varieties induced from the maps $\bar{\theta}$ and $\bar{\xi}$:

$$(4) \quad \widetilde{\text{Gr}}(q, V_{m,q}^*) \rightarrow \widetilde{\text{Gr}}(p, V_{n,p}^*) \text{ for } (n, p) \leq (m, q).$$

The above map is well defined since the following diagram is commutative:

$$(5) \quad \begin{array}{ccc} \widetilde{\text{Gr}}(p+1, V_{n+1,p+1}^*) & \xrightarrow{\bar{\xi}^{n+1,p+1}} & \widetilde{\text{Gr}}(p+1, V_{n,p+1}^*) \\ \downarrow \bar{\theta}^{n+1,p+1} & & \downarrow \bar{\theta}^{n,p+1} \\ \widetilde{\text{Gr}}(p, V_{n+1,p}^*) & \xrightarrow{\bar{\xi}^{n+1,p}} & \widetilde{\text{Gr}}(p, V_{n,p}^*). \end{array}$$

Define the affine schemes:

$$(6) \quad \widetilde{\text{Gr}}(\infty/2, V_\infty^*) := \varprojlim_{(n,p)} \widetilde{\text{Gr}}(p, V_{n,p}^*),$$

and

$$(7) \quad \wedge^{\infty/2}(V_\infty^*) := \varprojlim_{(n,p)} \wedge^p(V_{n,p}^*).$$

We call $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ the *affine infinite Grassmannian*.

Then, the corresponding affine coordinate rings are given by:

$$(8) \quad k[\widetilde{\text{Gr}}(\infty/2, V_\infty^*)] = \varinjlim_{(n,p)} k[\widetilde{\text{Gr}}(p, V_{n,p}^*)],$$

and

$$(9) \quad k[\wedge^{\infty/2}(V_\infty^*)] = \varinjlim_{(n,p)} k[\wedge^p(V_{n,p}^*)] = \varinjlim_{(n,p)} S^\bullet(\wedge^p(V_{n,p})),$$

where S^\bullet is the symmetric algebra.

The multiplicative group k^* acts on $\wedge^p(V_{n,p}^*)$ via multiplication by $z \in k^*$. Clearly, this k^* -action preserves $\widetilde{\text{Gr}}(p, V_{n,p}^*)$. Moreover, $\bar{\xi}$ and $\bar{\theta}$ both commute with this k^* -action. Thus, we get a k^* -action

on $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$. Further, $\widetilde{\xi}^{n+1, p+1}$ commutes with the standard $\text{GL}(V_{n, p+1})$ -actions (considering $\text{GL}(V_{n, p+1})$ as canonically embedded in $\text{GL}(V_{n+1, p+1})$) and $\widetilde{\theta}^{n+1, p+1}$ commutes with the standard $\text{GL}(V_{n+1, p})$ -actions (again considering $\text{GL}(V_{n+1, p})$ as canonically embedded in $\text{GL}(V_{n+1, p+1})$). In particular, $\widetilde{\xi}^{n+1, p+1}$ commutes with the $\text{N}(V_{n, p+1})$ -actions and $\widetilde{\theta}^{n+1, p+1}$ commutes with the $\text{N}(V_{n+1, p})$ -actions, where $\text{N}(V_{n+1, p})$ denotes the normalizer of the standard maximal torus in $\text{GL}(V_{n+1, p})$.

Define the group

$$\text{GL}(\infty) := \varinjlim_{(n, p)} \text{GL}(V_{n, p}),$$

and its subgroup

$$\text{N}(\infty) := \varinjlim_{(n, p)} \text{N}(V_{n, p}).$$

Then, $\text{GL}(\infty)$ acts on $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ as follows. Take $g \in \text{GL}(V_{n, p})$. Now,

$$(10) \quad \widetilde{\text{Gr}}(\infty/2, V_\infty^*) := \varprojlim_{(m, q) \geq (n, p)} \widetilde{\text{Gr}}(q, V_{m, q}^*).$$

Each variety on the right is acted upon by $\text{GL}(V_{n, p})$ and all the maps $\widetilde{\xi}$ and $\widetilde{\theta}$ are $\text{GL}(V_{n, p})$ -equivariant maps. Thus, $g \in \text{GL}(V_{n, p})$ acts on $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ for any pair (n, p) . These actions clearly combine to give an action of $\text{GL}(\infty)$ on $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$.

The standard action of $\text{GL}(V_{n, p})$ on $\wedge^p(V_{n, p}^*)$ commutes with the above k^* -action and hence so is the k^* -action on $\widetilde{\text{Gr}}(p, V_{n, p}^*)$ commutes with the standard $\text{GL}(V_{n, p})$ -action. Thus, we get a k^* -action on $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ commuting with the action of $\text{GL}(\infty)$.

Define a bijection

$$\mu : -\mathbb{N} \sqcup \mathbb{N} \rightarrow \mathbb{N}, \quad -n \mapsto 2n, \quad p \mapsto 2p - 1, \quad \text{for } n, p \in \mathbb{N},$$

where \mathbb{N} is the set of positive integers $\{1, 2, \dots\}$. This gives rise to a *well order* on $-\mathbb{N} \sqcup \mathbb{N}$ transporting the standard well order on \mathbb{N} via μ . Write a basis of $\wedge^p(V_{n, p})$ as follows:

$$x_{i_1} \wedge \cdots \wedge x_{i_p}, \quad \text{where } i_j \in \{-n, -(n-1), \dots, -1, 1, \dots, p\}$$

so that $i_1 < i_2 < \cdots < i_p$ in the above well order. Now, define

$$\mathbf{x}_i := x_{i_1} \wedge \cdots \wedge x_{i_p} < x_{j_1} \wedge \cdots \wedge x_{j_p}$$

under the lexicographic order reading from the left.

The following lemma is clear.

Lemma 2.2. *The above order on the basis of $\wedge^p(V_{n, p})$ is a well order.*

Lemma 2.3. *Under the embedding $\beta : \wedge^p(V_{n, p}) \hookrightarrow \wedge^{p+1}(V_{n, p+1})$, $\omega \mapsto \omega \wedge x_{p+1}$, the above well ordering on the basis of $\wedge^{p+1}(V_{n, p+1})$ restricts to the well ordering on the (up to signs) basis of $\wedge^p(V_{n, p})$.*

Proof. First, let $x_{i_1} \wedge \cdots \wedge x_{i_p} < x_{j_1} \wedge \cdots \wedge x_{j_p}$. Then, we claim that

$$(11) \quad x_{i_1} \wedge \cdots \wedge x_{i_p} \wedge x_{p+1} < x_{j_1} \wedge \cdots \wedge x_{j_p} \wedge x_{p+1} :$$

Choose the largest $\ell \geq 0$ such that $i_1 = j_1, \dots, i_\ell = j_\ell$. If $\mu(p+1) < \mu(i_\ell)$, then clearly the equation (11) is true. So, assume that $\mu(p+1) > \mu(i_\ell)$. If $\mu(p+1) > \mu(j_{\ell+1}) > \mu(i_{\ell+1})$, then again clearly the equation (11) is true. So, assume that $\mu(i_\ell) = \mu(j_\ell) < \mu(p+1) < \mu(j_{\ell+1})$. If

$\mu(i_{\ell+1}) > \mu(p+1)$, then again the equation (11) is true. So, finally assume that $\mu(i_{\ell+1}) < \mu(p+1)$ and $\mu(j_{\ell+1}) > \mu(p+1)$. Since $\mu(i_{\ell+1}) < \mu(p+1)$, the equation (11) is true since $\mu(j_{\ell}) < \mu(p+1) < \mu(j_{\ell+1})$.

Conversely, if

$$(12) \quad x_{i_1} \wedge \cdots \wedge x_{i_p} \wedge x_{p+1} < x_{j_1} \wedge \cdots \wedge x_{j_p} \wedge x_{p+1},$$

then we assert that $x_{i_1} \wedge \cdots \wedge x_{i_p} < x_{j_1} \wedge \cdots \wedge x_{j_p}$: For, otherwise, assume that $x_{i_1} \wedge \cdots \wedge x_{i_p} > x_{j_1} \wedge \cdots \wedge x_{j_p}$. This implies $x_{i_1} \wedge \cdots \wedge x_{i_p} \wedge x_{p+1} > x_{j_1} \wedge \cdots \wedge x_{j_p} \wedge x_{p+1}$, contradicting the equation (12). This proves the lemma. \square

The proof of the following lemma is clear.

Lemma 2.4. *Under the embedding $\eta : \wedge^p(V_{n,p}) \hookrightarrow \wedge^p(V_{n+1,p})$, $\omega \mapsto \omega$, the above well ordering on the basis of $\wedge^p(V_{n+1,p})$ restricts to the well ordering on the basis of $\wedge^p(V_{n,p})$.*

We recall the following definition from [AH, §3.3].

Definition 2.5. Let \leq be a well ordering on a countable set X . Define the induced *lexicographic ordering* \leq^o on the set X^o of commuting monomials with terms from X as follows:

$$(13) \quad \mathbf{x} := x_1^{m_1} \cdots x_a^{m_a} \leq^o \mathbf{y} := x_1^{n_1} \cdots x_a^{n_a} \Leftrightarrow (m_1, \dots, m_a) \leq (n_1, \dots, n_a)$$

lexicographically from the left, where $x_i \in X$, $x_1 < x_2 < \cdots < x_a$ and $m_i, n_i \in \mathbb{Z}_{\geq 0}$. Then, \leq^o is a term ordering on X^o (cf. [AH, Example2.5]). Let a group G act on X . Define a quasi-ordering $|_G$ on X^o by

$$\mathbf{x}|_G \mathbf{y} \Leftrightarrow \exists \sigma \in G \text{ and } \mathbf{z} \in X^o : (\sigma \mathbf{x})\mathbf{z} = \mathbf{y}.$$

Define the infinite symmetric group $S_\infty := \varinjlim S_n$, where S_n is the symmetric group on the symbols $\{-n, -(n-1), \dots, -1, 1, 2, \dots, n\}$. Then, S_∞ is canonically embedded as a subgroup of $\text{GL}(\infty)$ obtained via the permutation matrices.

Conjecture 2.6. *(due to J. Draisma) Let k be any field. The affine infinite Grassmannian $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ is topologically S_∞ -noetherian, i.e., every descending chain of S_∞ -stable Zariski-closed subsets of the k -points of $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ stabilizes.*

A slightly weaker form of the conjecture states that $\widetilde{\text{Gr}}(\infty/2, V_\infty^)$ is topologically $\text{N}(\infty)$ -noetherian.*

A stronger form of the conjecture states that any ascending chain of S_∞ -stable ideals in the affine coordinate ring of $\widetilde{\text{Gr}}(\infty/2, V_\infty^)$ stabilizes.*

3. A COUNTEREXAMPLE TO A STRONGER FORM OF THE MATROID MINOR CONJECTURE

We begin first by disproving the strong form of Matroid Minor Conjecture for $\wedge^{\infty/2}(V_\infty^*)$.

Proposition 3.1. *The ring*

$$R := k[\wedge^{\infty/2}(V_\infty^*)] = \varinjlim_{(n,p)} S^\bullet(\wedge^p(V_{n,p})) \quad (\text{cf. the identity (9)})$$

is not noetherian with respect to the S_∞ -stable ideals, i.e., there exists a strictly increasing sequence of S_∞ -stable ideals of R :

$$I_1 \subsetneq I_2 \subsetneq I_3 \subsetneq \dots$$

Proof. Using [AH, Lemma3.14], it suffices to show that $|_{S_\infty}$ is *not* well-quasi-ordering. Consider the monomials

$$(14) \quad \{a_n b_n\}_{n \geq 3}, \text{ where } a_n := x_{-2} \wedge x_2 \wedge x_3 \wedge \cdots \wedge x_n \\ \text{and } b_n := x_{-(n-1)} \wedge x_{-(n-2)} \wedge \cdots \wedge x_{-1} \wedge x_1 \wedge x_{n+1}.$$

We claim that $a_n b_n \not\leq_{S_\infty} a_m b_m$ for $n \neq m \geq 3$:

If not, let $\sigma \in S_\infty$ be such that $\sigma(a_n) \cdot \sigma(b_n) = a_m b_m$ (observe that they both are degree 2 monomials), i.e.,

$$\begin{aligned} & (x_{\sigma(-2)} \wedge x_{\sigma(2)} \wedge x_{\sigma(3)} \wedge \cdots \wedge x_{\sigma(n)}) \\ & \cdot (x_{\sigma(-(n-1))} \wedge x_{\sigma(-(n-2))} \wedge \cdots \wedge x_{\sigma(-1)} \wedge x_{\sigma(1)} \wedge x_{\sigma(n+1)}) \\ & = (x_{-2} \wedge x_2 \wedge x_3 \wedge \cdots \wedge x_m) \cdot (x_{-(m-1)} \wedge x_{-(m-2)} \wedge \cdots \wedge x_{-1} \wedge x_1 \wedge x_{m+1}). \end{aligned}$$

This gives

$$\begin{aligned} & \sigma(\{-2, 2, 3, \dots, n, n+1, \dots\}) \setminus \sigma(\{-(n-1), -(n-2), \dots, -1, 1, n+1, n+2, \dots\}) \\ & = \begin{cases} \{-2, 2, 3, \dots, m, m+1, \dots\} \setminus \{-(m-1), -(m-2), \dots, -1, 1, m+1, m+2, \dots\} \text{ or} \\ \{-(m-1), -(m-2), \dots, -1, 1, m+1, m+2, \dots\} \setminus \{-2, 2, 3, \dots, m, m+1, \dots\}. \end{cases} \end{aligned}$$

The above equation is equivalent to the following:

$$\sigma(\{2, 3, \dots, n\}) = \begin{cases} \{2, 3, \dots, m\} \text{ or} \\ \{-(m-1), -(m-2), \dots, -3, -1, 1\}. \end{cases}$$

The left side of the above equation has cardinality $n-1$, whereas the right side has cardinality $m-1$. This is a contradiction since $n \neq m$ by assumption. Thus, the infinite set $A = \{a_n b_n\}_{n \geq 3}$ is an anti-chain under $|_{S_\infty}$ (cf. [AH, Page 5173]). This proves the proposition. \square

Remark 3.2. *The ring R as in Proposition 3.1 is not noetherian even with respect to $\text{GL}(\infty)$ -stable ideals as observed in [DE, Open Problem 1].*

Consider the ordered basis

$$x_{-1}, x_{-2}, \dots, x_{-(m-1)}, x_1, x_2, \dots, x_{m+1} \text{ of } V_{m-1, m+1}.$$

We abbreviate $m+1$ by p . Consider the standard maximal parabolic subgroup $P(m-1)$ of $\text{SL}(2m)$ (with respect to the above ordered basis) obtained by deleting the $(m-1)$ -th node from the Dynkin diagram of $\text{SL}(2m)$. Thus, $P = P(m-1)$ is the stabilizer of the line $[x_1^* \wedge x_2^* \wedge \cdots \wedge x_p^*] \in \mathbb{P}(\wedge^p(V_{m-1, p}^*))$. Consider the opposite unipotent radical $U^- = U_{m-1}^-$ of P . Thus, for any $g \in U^-$,

$$\begin{aligned} g^{-1}(x_{-j}) &= x_{-j} + \sum_{i=1}^p \alpha_i^j(g) x_i, \text{ for } 1 \leq j \leq m-1, \text{ and} \\ g^{-1}(x_i) &= x_i, \text{ for } 1 \leq i \leq p, \end{aligned}$$

for some $\alpha_i^j(g) \in k$. In fact, U^- is characterized by the above, where we allow $\alpha_i^j(g)$ to vary over k . The action of U^- on the dual basis is given as follows (for $g \in U^-$):

$$\begin{aligned} g(x_{-j}^*) &= x_{-j}^*, \text{ for } 1 \leq j \leq m-1, \text{ and} \\ g(x_i^*) &= x_i^* + \sum_{j=1}^{m-1} \alpha_i^j(g) x_{-j}^*, \text{ for } 1 \leq i \leq p, \end{aligned}$$

Thus, for $g \in U^-$,

$$(15) \quad g(x_1^* \wedge x_2^* \wedge \cdots \wedge x_p^*) = \left(x_1^* + \sum_{j_1=1}^{m-1} \alpha_1^{j_1}(g)x_{-j_1}^* \right) \wedge \cdots \wedge \left(x_p^* + \sum_{j_p=1}^{m-1} \alpha_p^{j_p}(g)x_{-j_p}^* \right).$$

With the notation as above, we have the following lemma.

Lemma 3.3. *For $g \in U^-$ and $\mathbf{x} = x_{-n_1} \wedge \cdots \wedge x_{-n_q} \wedge x_{d_1} \wedge \cdots \wedge x_{d_{p-q}}$, where $0 \leq q \leq m-1$, $0 < n_1 < \cdots < n_q \leq m-1$ and $0 < d_1 < \cdots < d_{p-q} \leq p$, we have the following identity:*

$$\begin{aligned} & g(x_1^* \wedge x_2^* \wedge \cdots \wedge x_p^*) (x_{-n_1} \wedge \cdots \wedge x_{-n_q} \wedge x_{d_1} \wedge \cdots \wedge x_{d_{p-q}}) \\ &= \pm \det(\alpha_{m_j}^{n_i})_{1 \leq i \leq q; m_j \in \{1, 2, \dots, \widehat{d_1}, \dots, \widehat{d_{p-q}}, \dots, p\}}. \end{aligned}$$

Proof.

$$g(x_1^* \wedge x_2^* \wedge \cdots \wedge x_p^*) \mathbf{x} = \det \begin{pmatrix} \alpha_1^{n_1}(g) & \cdots & \alpha_1^{n_q}(g) & 0 & 0 & \cdots & 0 \\ \alpha_2^{n_1}(g) & \cdots & \alpha_2^{n_q}(g) & 0 & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \alpha_{d_1}^{n_1}(g) & \cdots & \alpha_{d_1}^{n_q}(g) & 1 & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \alpha_{d_2}^{n_1}(g) & \cdots & \alpha_{d_2}^{n_q}(g) & 0 & 1 & \cdots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \alpha_{d_{p-q}}^{n_1}(g) & \cdots & \alpha_{d_{p-q}}^{n_q}(g) & 0 & 0 & \cdots & 1 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ \alpha_p^{n_1}(g) & \cdots & \alpha_p^{n_q}(g) & 0 & 0 & \cdots & 0 \end{pmatrix}.$$

By moving d_1, d_2, \dots, d_{p-q} -th rows of the above matrix to the end of the matrix, we get (from the above equation),

$$g(x_1^* \wedge x_2^* \wedge \cdots \wedge x_p^*) \mathbf{x} = \pm \det(\alpha_{m_j}^{n_i})_{1 \leq i \leq q; m_j \in \{1, 2, \dots, \widehat{d_1}, \dots, \widehat{d_{p-q}}, \dots, p\}}.$$

This proves the lemma. \square

Let $\widetilde{\text{Gr}}(\infty/2, V_\infty^*) \hookrightarrow \wedge^{\infty/2}(V_\infty^*)$ be the Plücker embedding induced from the Plücker embeddings $\widetilde{\text{Gr}}(p, V_{n,p}^*) \hookrightarrow \wedge^p(V_{n,p}^*)$ and let $\mathcal{I} \subset k[\wedge^{\infty/2}(V_\infty^*)]$ be the ideal generated by the Plücker relations, i.e., the ideal of the subscheme $\widetilde{\text{Gr}}(\infty/2, V_\infty^*)$ embedded in $\wedge^{\infty/2}(V_\infty^*)$ via the Plücker embedding (cf. [EH, §III.2.7]). Consider the sequence of ideals in $k[\wedge^{\infty/2}(V_\infty^*)]$:

$$(16) \quad \mathcal{I}_n := \langle S_\infty(a_3 \cdot b_3), \dots, S_\infty(a_n \cdot b_n) \rangle + \mathcal{I},$$

where a_i, b_i are defined by the equation (14) and $S_\infty(a_i \cdot b_i)$ denotes the collection $\{\sigma(a_i) \cdot \sigma(b_i)\}_{\sigma \in S_\infty}$.

We have the following main theorem of this note.

Theorem 3.4. *The above ideals satisfy:*

$$\mathcal{I}_3 \subsetneq \mathcal{I}_4 \subsetneq \mathcal{I}_5 \subsetneq \dots$$

In particular, the ring $\mathcal{R} := k[\widetilde{\text{Gr}}(\infty/2, V_\infty^)]$ is not noetherian with respect to the S_∞ -stable ideals. Thus, the stronger form of the Matroid Minor Conjecture (cf. Conjecture 2.6) is false.*

In fact, as mentioned by A. Snowden, the ideals \mathcal{I}_n are even stable under $N(\infty)$ since the $a_i b_i$ are eigenvectors for the action of the maximal torus.

Proof. Fix $\ell > 3$. It suffices to show that $a_\ell \cdot b_\ell \notin \mathcal{I}_n$ for any $n < \ell$. This is equivalent to proving that for any $\sigma_i \in k[S_\infty]$, where $k[S_\infty]$ is the group algebra of S_∞ ,

$$a_\ell \cdot b_\ell - \sum_{i=3}^{\ell-1} \sigma_i(a_i \cdot b_i) \text{ does not vanish identically on } \widetilde{\text{Gr}}(\infty/2, V_\infty^*).$$

Write

$$\sigma_i = \sum_k z_i^k \sigma_i^k \text{ (a finite sum) for some } z_i^k \in k \text{ and } \sigma_i^k \in S_\infty.$$

Take $m \geq \ell$ large enough so that each σ_i^k (with nonzero z_i^k) is a permutation of the basis of $V_{m-1,p:=m+1}$. We want to show that

$$(17) \quad a_\ell \cdot b_\ell - \sum_{i=3}^{\ell-1} \sum_k z_i^k \sigma_i^k(a_i \cdot b_i) \text{ thought of as a function on } \wedge^p(V_{m-1,p}^*) \text{ does not vanish identically on } U_m^-.$$

Now,

$$\begin{array}{ccc} S^\bullet(\wedge^p(V_{m-1,p})) & \hookrightarrow & k[\wedge^{\infty/2}(V_\infty^*) \\ \downarrow & & \downarrow \\ k[\widetilde{\text{Gr}}(p, V_{m-1,p}^*)] & \hookrightarrow & k[\widetilde{\text{Gr}}(\infty/2, V_\infty^*)], \end{array}$$

where S^\bullet denotes the symmetric algebra and both the vertical maps are surjective. Following the notation in Lemma 3.3, we rewrite

$$\det(\alpha_{m_j}^{n_i})_{1 \leq i \leq q; m_j \in \{1, 2, \dots, \widehat{d_1}, \dots, \widehat{d_{p-q}}, \dots, p\}} = \det(\alpha_{m_1, \dots, m_q}^{n_1, \dots, n_q}).$$

Assume, if possible, that

$$(18) \quad \left(g(x_1^* \wedge x_2^* \wedge \dots \wedge x_p^*) \right) \left(a_\ell \cdot b_\ell - \sum_{i=3}^{\ell-1} \sum_k z_i^k \sigma_i^k(a_i \cdot b_i) \right) = 0, \text{ for all } g \in U_m^-.$$

By Lemma 3.3, we get

$$g(x_1^* \wedge x_2^* \wedge \dots \wedge x_p^*)(a_\ell \cdot b_\ell) = \alpha_1^2 \det(\alpha_{2,3,\dots,\ell}^{1,\dots,\ell-1}).$$

Now, if $g(x_1^* \wedge x_2^* \wedge \dots \wedge x_p^*)(\sigma_i^k(a_i \cdot b_i))$ has any nonzero contribution to the above term, considering the action of k^* on each of the variables α_j^i , for a fixed i (and any j) and similarly for k^* -action on α_j^i for a fixed j (and any i) by the same character, we should have (by Lemma 3.3):

$$\sigma_i^k(A) = \{-C, -2, \ell + 1, \dots, p, C'\},$$

where $A := \{-2, 2, 3, \dots, p\}$ and

$$\sigma_i^k(B) = \{-D, -2, \ell + 1, \dots, p, D'\},$$

for some C, C', D, D' satisfying the following:

$$\{1, \widehat{2}, 3, \dots, \ell - 1\} = C \sqcup D \text{ and } \{1, 2, 3, \dots, \ell\} = C' \sqcup D',$$

where $B := \{-(i-1), -(i-2), \dots, -1, 1, i+1, i+2, \dots, p\}$.

Setting $c = |C|$, we get

$$|D| = \ell - 2 - c, |C'| = -(p - \ell + c + 1) + p = \ell - c - 1, \text{ and } |D'| = -(p - \ell + 1 + \ell - 2 - c) + p = c + 1.$$

Then,

$$(19) \quad \sigma_i^k(A \setminus B) = \sigma_i^k(\{2, 3, \dots, i\})$$

and

$$(20) \quad \sigma_i^k(B \setminus A) = \sigma_i^k(\{-(i-1), -(i-2), \dots, -3, -1, 1\}).$$

But,

$$(21) \quad \sigma_i^k(A) \setminus \sigma_i^k(B) = \{-C, C'\}$$

and

$$(22) \quad \sigma_i^k(B) \setminus \sigma_i^k(A) = \{-D, D'\}.$$

This leads to a contradiction for any $i < \ell$, since (by the equations (19) and (21)),

$$|\sigma_i^k(A \setminus B)| = i - 1 = \ell - 1.$$

Also, by the equations (20) and (22),

$$|\sigma_i^k(B \setminus A)| = i - 1 = \ell - 1.$$

This contradiction proves that the equation (18) cannot be true. This proves the theorem. \square

REFERENCES

- [AH] M. Aschenbrenner and C. Hillar, *Finite generation of symmetric ideals*. *Transactions of A.M.S.* **359** (2007), 5171–5192.
- [Di] R. Diestel, *Graph theory*, Springer-Verlag, New York, 1997.
- [DE] J. Draisma and R. H. Eggermont, Plücker varieties and higher secants of Sato’s Grassmannian, *J. Reine Angew. Math.* **737** (2018), 189–215.
- [EH] D. Eisenbud and J. Harris, *The Geometry of Schemes*, GTM volume 197, Springer (2000).
- [GGW] J. Geelen, B. Gerards and G. Whittle, Solving Rota’s conjecture, *Notices of the AMS* **61**, Number 7, (August 2014), 736–743.
- [Kur] K. Kuratowski, Sur le problème des courbes gauches en topologie, *Fund. Math.* **15** (1930), 271–283.
- [RS] N. Robertson and P. D. Seymour, Graph Minors. XX. Wagners Conjecture, *J. Combin. Theory Ser. B* **92** (2004), 325–357.
- [Ro] G.-C. Rota, Combinatorial theory, old and new, in: *Proc. Internat. Cong. Math. (Nice, 1970)*, pp. 229–233. Gauthier-Villars, Paris.

S. KUMAR: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27599-3250, USA
E-mail address: shrawan@email.unc.edu